Objects That Cannot Be Taken Apart With Two Hands

Jack Snoeyink*
Department of Computer Science
University of British Columbia

Jorge Stolfi†
Department of Computer Science
University of Campinas

Abstract

It has been conjectured that every configuration $\mathcal{C}$ of convex objects in 3-space with disjoint interiors can be taken apart by translation with two hands: that is, some proper subset of $\mathcal{C}$ can be translated to infinity without disturbing its complement. We show that the conjecture holds for five or fewer objects and give a counterexample with six objects. We extend the counterexample to a configuration that cannot be taken apart with two hands using arbitrary isometries (rigid motions).

1 Introduction

One can define an *aboveness* relation for convex objects in a Euclidean space E^d by saying that A is *above* B if a vertical ray drawn upward from a point in the interior of B encounters A. (Objects in this paper always have disjoint interiors.) In the plane, this relation is acyclic [3, 4]—a fact that is useful in many algorithms [5, 10]. This also implies that there is a maximum object with respect to aboveness. Thus, a planar configuration $\mathcal{C}$ can be taken apart (or assembled if we reverse time) by repeatedly translating a maximum object upwards to infinity.

*Supported in part by an NSERC Research Grant

†Previously of DEC Systems Reseach Center, Palo Alto, CA

Computer graphics textbooks often illustrate that this property does not hold for objects in 3-space: three disjoint convex objects can overlap cyclically for the correct choice of "upward." In 1984, Dawson [2] showed an example of 12 convex tiles, none of which can be translated "upward" without disturbing the others for any choice of "upward." In 1988, Natarajan [8] gave an example with 16 thin triangular plates. Already in 1963, Fejes-Toth and Heppes [4] had given an example of 13 convex objects, 12 tetrahedra surrounding a rhombic dodecahedron, for which no single object can move rigidly (according to an arbitrary isometry) without disturbing the others. In section 4 we confirm their conjecture that the central dodecahedron is not necessary.

Because all three examples essentially tile a sphere with tiles whose neighbors allow movement only in the plane of the tile, Natarajan [8] conjectured that every configuration $\mathcal{C}$ of convex objects could be *taken apart by translation with two hands:* that there always exists a proper subset $\mathcal{A} \subset \mathcal{C}$ that can be translated to infinity by applying a common translation to the objects in $\mathcal{A}$, without disturbing the objects in the complement $\mathcal{C} \setminus \mathcal{A}$. (No point of $\mathcal{A}$ enters the interior of an object in $\mathcal{C} \setminus \mathcal{A}$ during the translation.)

In this paper, we construct counterexamples to this conjecture. Section 2 gives a configuration of six tetrahedra and a proof that it cannot be taken apart by translation with two hands. The construction is based on the alternating group A^4. Section 3 shows that this configuration is minimal, in the sense that one object out of any configuration

of five convex objects can always be translated to infinity. Section 4 turns to arbitrary isometries and confirms a conjecture that Fejes-Toth and Heppes made about their construction, and section 5 gives a configuration of 30 objects that cannot be taken apart by applying an isometry to any proper subset. The 30 objects are formed by lacing five copies of the translational counterexample of section 2 together, based on five subgroups of A^4 that can be found in the alternating group A^5. The proof of this case is an exhaustive computer check of all non-symmetric cases, using an analysis technique for positive-force closure grasps in robotics [7, 9] to check for possible infinitesimal rigid motions.

De Bruijn [3], and later Dawson [2], proved that any collection of n star-shaped objects with disjoint interiors can be separated by n hands using translation: For each object, choose a point q_i from which object i is star-shaped, then translate object i with the velocity vector $q_i - q_1$. Since this translation could be accomplished by scaling the entire configuration about q_1 and then shrinking each object i radially about q_i to its original size, there are no intersections during translation. Therefore, n hands are sufficient for convex objects. Our proof in the next section shows that three hands are necessary for a configuration of six objects. It would be interesting to narrow this gap.

2 A twisted tetrahedron

In this section we construct a configuration of six identical convex sticks that cannot be taken apart with two hands using translational motions. Our construction, illustrated in figure 1, essentially extends the edges of a tetrahedron and gives them a clockwise twist around each vertex. Building our configuration to have the symmetries of a regular tetrahedron reduces the number of subsets that must be checked for possible disassembly.

Consider a regular tetrahedron centered at the origin having vertices $\vec{a} = (+1,+1,+1)$, $\vec{b} = (+1,-1,-1)$, $\vec{c} = (-1,+1,-1)$, and $\vec{d} = (-1,-1,+1)$. The group of symmetries of this tetrahedron is the alternating group A^4, which can be generated by two rotation matricies:

$$A^4 = \{N, R \mid N^3 = I, R^2 = I, (NR)^3 = I\},$$

where

$$R = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & +1 \end{pmatrix}, N = \begin{pmatrix} 0 & +1 & 0 \\ 0 & 0 & +1 \\ +1 & 0 & 0 \end{pmatrix}.$$

We begin our construction with the 12 points formed by A^4 acting on the point $p = (6, 5, -2)$. Because the line segment joining p and $pR = (-6, -5, -2)$ is invariant under rotation R, the orbit of segment $(p; pR)$ consists of six different segments, each generated twice. We then define planes separating nearby segments—the six objects shown in figure 1 are the intersections of halfspaces defined by separating planes and containing the six segments.

We prove four properties of our construction: First, the intersections of halfplanes define non-empty, bounded, convex objects. Second, these objects have disjoint interiors. Third, they share 2-dimensional contact regions. Fourth, for any proper subset of these objects, every possible translation causes a moving object to penetrate a stationary object at some contact region.

The three points p, $pN = (-2, 6, 5)$, and pR define a plane equation $92 + \vec{n} \cdot q = 0$, where $\vec{n} = (35, -42, 46)$ is the normal vector. A point q is said to be in the *non-positive halfspace* of the plane if $92 + \vec{n} \cdot q \leq 0$. Notice that symmetries of the tetrahedron, which are pure rotations, act on the normal vector as they do on points and do not change the constant in a plane equation. Thus, we can name the 12 planes in the orbit of $92 + \vec{n} \cdot q = 0$ by their normal vectors. Four planes in the orbit contain either p or pR (see table 1.) We can orient these planes so that plugging in the point $q = (0, 0, -5/2)$, for example, gives negative values for all four planes. We call the intersection of the four non-positive halfspaces the *stick* Ξ.

Lemma 1 *The stick Ξ, which is the intersection of the non-positive halfspaces of $92 + \vec{n} \cdot q$, $92 + \vec{n}R \cdot q$, $-92 - \vec{n}NN \cdot q$ and $-92 - \vec{n}NNR \cdot q$, is a non-empty, bounded tetrahedron.*

Proof: The region is non-empty: $(0, 0, -5/2)$ is in the interior of the non-positive halfspaces of the four planes.

The region is also bounded: because the positive linear combination of the normals $35(\vec{n} + \vec{n}R) +$



Figure 1: Three views of six objects that cannot be taken apart by two hands using translations

Points defining a plane of Ξ	Normal vector	Plane eqn. LHS
$\{p,\ pN,\ pR\}$	$\vec{n} = (35, -42, 46)$	$92 + \vec{n} \cdot q$
$\{pR,\ (pN)R,\ (pR)R = p\}$	$\vec{n}R = (-35, 42, 46)$	$92 + \vec{n}R \cdot q$
$\{pNN,\ (pN)NN = p,\ (pR)NN\}$	$-\vec{n}NN = (42, -46, -35)$	$-92 - \vec{n}NN \cdot q$
$\{pNNR, (pN)NNR = pR, (pR)NNR\}$	$-\vec{n}NNR = (-42, 46, -35)$	$-92 - \vec{n}NNR \cdot q$

Table 1: The four planes that contain p or pR and define stick Ξ.

$46(-\vec{n}NN - \vec{n}NNR) = 0$, every vector in the subspace spanned by the normals can be written as a positive linear combination of the normals. One can verify that the normals are not coplanar, so every vector has positive projection on the normal to some plane, and motion along the vector will eventually leave the non-positive halfspace of that plane.

The only non-empty, bounded region defined by four planes is a tetrahedron. ■

We can apply the group action on Ξ to obtain an orbit with six different sticks, each generated twice.

Lemma 2 *The six sticks defined by group action of A^4 on Ξ have disjoint interiors.*

Proof: Stick ΞN uses the plane with equation $-92 - (\vec{n}NN)N \cdot q = -92 - \vec{n} \cdot q$. Since Ξ uses this plane with the opposite sign, the interiors of ΞN and Ξ are separated. In the same way, the interiors of ΞNR, ΞNN, and ΞNNR are separated from that of Ξ.

The only remaining stick is $\Xi NRNN$, which in figure 1 is clearly disjoint from Ξ. For a numerical proof, one can check that the plane $(0, 0, 1) \cdot q = 0$ separates the vertices of the two sticks. ■

Next, we observe that sticks that share a plane have a two-dimensional contact. We call such sticks *neighbors.* Because of symmetries, it is sufficient to check that Ξ and ΞN are neighbors.

Lemma 3 *The intersection of sticks Ξ and ΞN is a two-dimensional region.*

Proof: The point $r = (-3404, -3105, -2975)/1365$ satisfies $92 + \vec{n} \cdot r = 0$, so r is in the plane shared by sticks Ξ and ΞN. The point r is strictly below the unshared planes of these two sticks—plugging r into the left-hand sides of the plane equations gives strictly negative numbers—therefore r is contained in a two-dimensional polygon $\Xi \cap \Xi N$. ■

Theorem 4 *Figure 1 illustrates a configuration $\mathcal{C}$ of six tetrahedra that cannot be taken apart by translation with two hands.*

Proof: A stationary point in contact with the interior of a face of a moving object rules out all translations with positive projection onto the normal of that face. To show that a proper subset of $\mathcal{C}$ cannot translate infinitesimally, it is enough to show that any direction has positive projection on such a contact normal.

We consider all possible subsets, up to symmetry. Whether a subset or its complement moves depends on one's point of view, so we can focus on subsets of at most three sticks.

The proof of lemma 1 has already shown that every vector can be expressed as a positive linear combination of the normals of a single stick. Thus, no stick with four stationary neighbors can translate.

Next, suppose that a moving subset $\mathcal{A}$ contains a stick Ξ that has three stationary neighbors. We show that $\mathcal{A}$ can translate only if $\mathcal{A}-\{\Xi\}$ can translate. Let $\vec{n}$ be the normal of Ξ at the contact with its moving neighbor and let $\vec{n}_1$, $\vec{n}_2$, and $\vec{n}_3$ be the normals at stationary contacts. The vector $-\vec{n}$ can be expressed as a positive linear combination of $\vec{n}_1$, $\vec{n}_2$, and $\vec{n}_3$, by lemma 1. If we treat Ξ as stationary and find that some translation $\vec{t}$ is ruled out for $\mathcal{A}-\{\Xi\}$ because $(-\vec{n})\cdot\vec{t}>0$, then $\vec{n}_i\cdot\vec{t}>0$ for some $\vec{n}_i$ and translation $\vec{t}$ is also ruled out for $\mathcal{A}$ by motion of Ξ into one of its stationary contacts.

The only possible subsets remaining are three sticks that form a "tripod" or a "triangle"—these subsets are complements. We therefore consider the tripod formed by sticks Ξ, ΞN, and ΞNN. The normals at stationary contacts are

Stick	Contact normal
Ξ :	$\vec{n}R = (-35,\ 42,\ 46)$
	$-\vec{n}NNR = (-42,\ 46,-35)$
ΞN :	$\vec{n}RN = (\ 46,-35,\ 42)$
	$-\vec{n}NNRN = (-35,-42,\ 46)$
ΞNN :	$\vec{n}RNN = (\ 42,\ 46,-35)$
	$-\vec{n}NNRNN = (\ 46,-35,-42)$

These normals span the space of all vectors and
$31(\vec{n}R+\vec{n}RN+\vec{n}RNN)$
$+53(-\vec{n}NNR-\vec{n}NNRN-\vec{n}NNRNN)=0$.

Therefore, we conclude that no proper subset can be translated without penetrating its stationary complement. ■

3 One of five can translate

To prove that the construction of the previous section is minimal, we show first that any member of any configuration $\mathcal{C}$ of four or fewer convex objects can be translated to infinity and second that some member of any configuration $\mathcal{C}$ of five convex objects can be translated to infinity.

We call a set of planes Π a *separating set* for an object $A\in\mathcal{C}$ if, for any object $B\in\mathcal{C}$, $B\neq A$, some plane $\pi_{AB}\in\Pi$ separates the interiors of A and B. The set Π is *minimal* if no proper subset of Π is a separating set for A. The *region of* A with respect to a separating set Π is the intersection of halfspaces defined by planes of Π that contain A. We observe that objects that can be translated to infinity have unbounded regions.

Lemma 5 *A member A of a set $\mathcal{C}$ of convex objects with disjoint interiors can be translated to infinity if and only if the region of A is unbounded w.r.t. some separating set of planes.*

The 'if' part of this lemma has an easy corollary:

Corollary 6 *Any member of a configuration $\mathcal{C}$ of at most four convex objects can be translated to infinity in some direction.*

Proof: A minimal separating set for an object $A\in\mathcal{C}$ consists of at most three planes, which divide 3-space into eight unbounded regions. ■

With difficulty that is disproportionate to the importance of the theorem, we can prove

Theorem 7 *Some member of a configuration $\mathcal{C}$ of five convex objects can be translated to infinity.*

We prove this theorem by supposing that there is a counterexample consisting of five disjoint objects and then deriving a contradiction. First we prove that the objects can be assumed to be bounded tetrahedra defined by ten separating planes and that any pair have intersecting faces. We will then choose a set P of ten points in these intersection regions: one point common to each pair of tetrahedra. We study the structure of the convex hull $CH(P)$, and prove that it has a ring of five triangular faces on its boundary. Two adjacent triangles must be on the same "side" of this ring; we can use those triangles to show that the objects are not disjoint.

Lemma 8 *If there is a counterexample to theorem 7 then there is one with five bounded tetrahedra A, B, C, D, and E defined by the ten planes π_{AB}, π_{AC}, $\ldots\pi_{DE}$ that separate pairs. Furthermore, faces defined in the same plane intersect: e.g. $A\cap B$ is a polygon in π_{AB}.*

Proof: Suppose we are given a configuration $\mathcal{C}$ of five convex objects with disjoint interiors such that no object can be translated to infinity; we construct a configuration of tetrahedra. Choose a minimal set of planes Π that is a separating set for each of the objects. Note that Π contains at most ten planes: one plane π_{AB} for each pair of objects $A, B \in \mathcal{C}$. If Π contains fewer than ten planes then some object $X \in \mathcal{C}$ has a separating subset of at most three planes; the region of X is unbounded and lemma 5 states that X can be translated to infinity. We therefore assume that every object in $\mathcal{C}$ has a separating subset of four planes of Π.

Since four planes in 3-space define only one bounded region, a tetrahedron, we can further assume that each object is contained in the tetrahedron defined by its four separating planes. We can expand each object to fill its tetrahedral region: Π remains a separating set for the expanded objects and translation becomes only more difficult.

For each pair of expanded objects (tetrahedra) A and B, we show that the intersection $A \cap B$ is a two-dimensional region in the separating plane π_{AB}.



Figure 2: Faces in π_{AB}

Initially, let π equal π_{AB}. If the interiors of the faces $A \cap \pi$ and $B \cap \pi$ do not intersect, then these faces can be separated in π by a line ℓ as in figure 2. Rotate π around ℓ so that π continues to separate A and B, until π hits the vertex of A or of B that is not in plane π_{AB}. Suppose, without loss of generality, that π hits the vertex of A. Then the set $\{\pi, \pi_{AC}, \pi_{AD}, \pi_{AE}\}$ is a separating set of planes for A that all pass through a common point. In this set, the region of A must be unbounded and A can translate to infinity. ■

We denote the vertex of the tetrahedron A that is $\pi_{AC} \cap \pi_{AD} \cap \pi_{AE}$ by A_B because it is opposite the face that intersects B. Let V denote the set of all twenty vertices from the five tetrahedra.

As illustrated in figure 3, the *A_B–cone* is the region defined by π_{AC}, π_{AD}, and π_{AE} that contains A; the *A_B–anticone* is the region separated from A by π_{AC}, π_{AD}, and π_{AE}. There are three facts to notice about cones and anticones. First, any plane that cuts the A_B–cone in a triangle does not intersect the A_B–anticone, and vice versa. Second, A will be strictly inside the convex hull of any four points chosen one from the interior of each of the four anticones defined by A's vertices. And third, if the vertex A_B is on the convex hull $CH(V)$, then the A_B–anticone is entirely outside $CH(V)$.

Because tetrahedra A and B intersect in a 2-dimensional region in π_{AB}, we can choose a point $p_{AB} \in A \cap B$ that lies on no planes of $\Pi \setminus \{\pi_{AB}\}$. For each successive (unordered) pair of objects X and Y, we can choose p_{XY} in $X \cap Y$ such that the only plane of Π containing p_{XY} is π_{XY} and p_{XY} is in general position with respect to the previously chosen points. (No three points collinear, no four points coplanar, and no three defining a plane parallel to a separating plane.) Let P denote the set of all ten points.



Figure 3: A_B–cone and A_B–anticone

Lemma 9 *Suppose that the A_B–anticone does not intersect $CH(P)$. Then the points p_{CD}, p_{DE} and p_{CE} form a face of $CH(P)$ that lies on the B side of π_{AB}, is contained in the B_A–anticone, and whose plane cuts the B_A–anticone in a triangle.*

Proof: Let Δ denote the triangle defined by p_{CD}, p_{DE} and p_{CE} and let γ denote the plane containing Δ. By convexity, the edges of Δ are contained in tetrahedra C, D, and E.

By assumption, p_{CD} is not in the A_B–anticone. Since planes π_{AC} and π_{AD} separate A and p_{CD}, we know that the entire region containing p_{CD} that is defined by π_{AC} and π_{AD} is on the A-side of π_{AE}. Therefore, π_{AE} cuts through the triangle Δ and separates this region from E. Similarly, planes π_{AC} and π_{AD} cut through Δ and Δ cuts the A_B–cone in a triangle.

Now, we know that rays from the point p_{CD} through E intersect planes π_{AE}, π_{CE}, and π_{DE}

before E. By lemma 5 they must intersect π_{BE} afterwards (otherwise E would be in an unbounded region). Similar arguments apply to p_{CE} and D and to p_{DE} and C. Thus, in the plane γ, the planes π_{BC}, π_{BD}, and π_{BE} form a triangle that contain p_{CD}, p_{DE} and p_{CE}. In other words, the plane γ cuts the B_A–anticone in a triangle that contains Δ.

Since B is a bounded tetrahedron defined by planes π_{AB}, π_{BC}, π_{BD}, and π_{BE}, there is no point in space that separated from B by all four planes. Therefore, Δ is on the B side of π_{AB}.

Because the plane γ cuts the B_A–anticone in a triangle, γ does not intersect B. Moreover, γ cuts the A_B–cone on the B side of the separating plane π_{AB}. Therefore γ does not intersect A. Point p_{AB} is common to A and B, so both A and B lie strictly to one side of γ. Since all other points of P lie on A or B, γ certifies that p_{CD}, p_{DE} and p_{CE} form a face of $CH(P)$. ■

As a corollary of this lemma, we can show that if no points of $CH(P)$ are in the A_B–anticone, then there must be points of $CH(P)$ in several other anticones. For example, at most one anticone from each tetrahedron is free of points of $CH(P)$.

Corollary 10 *If the A_B–anticone is free of points of $CH(P)$ then the B_A–anticone, other anticones from A, and other anticones opposite B must contain points of P.*

Proof: Suppose the A_B–anticone is free of points of $CH(P)$. Then lemma 9 says that $p_{CD} \in CH(P)$ is contained in the B_A–anticone.

Lemma 9 also says that p_{CD} is on the E side of π_{BE}, since it is the B_A–anticone. Therefore, p_{CD} cannot be in the E_A–anticone and the A_E–anticone must contain points of $CH(P)$. Similarly, p_{CE} is on the D side of π_{BD} and p_{DE} is on the C side of π_{BC}, so the A_D–anticone and A_C–anticone also contain points of $CH(P)$.

Finally, lemma 9 says that p_{CD} is on the B side of AB. Therefore, p_{CD} cannot be in the B_C–anticone and the C_B–anticone contains points of P. Similar arguments apply for the D_B–anticone and E_B–anticone. ■

From this we can determine that all ten points of P appear on their convex hull, $CH(P)$.

Lemma 11 *By relabeling tetrahedra, we can assume that the following five triangles appear as faces of $CH(P)$: $\triangle p_{CD}p_{DE}p_{CE}$, $\triangle p_{AD}p_{DE}p_{AE}$, $\triangle p_{AB}p_{BE}p_{AE}$, $\triangle p_{AB}p_{BC}p_{AC}$, and $\triangle p_{BC}p_{CD}p_{BD}$.*

Proof: We show that each tetrahedron has an anticone that is free of points of P, then argue that corollary 10 permits only one listing of anticones, up to relabeling.

Notice that if any tetrahedron has points of P in each of its anticones, then that tetrahedron is strictly inside the convex hull $CH(P)$. Now, the convex hull $CH(V)$ has a vertex, which we can label A_B. Because A_B–anticone does not intersect $CH(P)$, lemma 9 says that points p_{CD} and p_{DE} lie on $CH(P)$. But this implies that the tetrahedra C, D, and E are not strictly inside $CH(P)$—they each must have an anticone free of points of P. The corollary implies that these anticones cannot be opposite B, so lemma 9 applied to one of these anticones will show that a point of B also lies on $CH(P)$—B also has an empty anticone.

Corollary 10 shows that there is one anticone per tetrahedron and one of each subscript. With five anticones, this can only occur as a five-cycle A_B, B_C, C_D, D_E, and A_E or as a three- and two-cycle. Since two-cycles (e.g. A_B and B_A) are explicitly ruled out, we must have the five-cycle, up to relabeling. Applying lemma 9 to this cycle gives the result. ■



Figure 4: The ring of faces

We can now finish the proof of theorem 7 by showing that two objects must intersect. As figure 4 illustrates, the five triangular faces found by lemma 11 form a ring on the convex hull $CH(P)$.

Since the ring has odd length, there must be two adjacent triangles that lie on the same side of the ring: they are $\triangle p_{CD}p_{DE}p_{CE}$ and $\triangle p_{AD}p_{DE}p_{AE}$ in the figure. But then the edges incident on the common vertex alternate: coming from objects D, E, D, E in the figure. Because the objects are convex, this alternation means that the interiors of D and E are not disjoint. This contradicts our initial assumption, and establishes the theorem. ■

4 The example of Fejes-Toth and Heppes

After seeing that not all configurations can be taken apart with two hands using translation, one naturally asks about other rigid motions. As mentioned in the introduction, Fejes-Toth and Heppes [4] have given a configuration of 13 objects—12 tetrahedra packed around a rhombic dodecahedron—in which no single object can move by any infinitesimal rigid motion (isometry). They also conjectured that the central dodecahedron was unnecessary. In this section, we prove this conjecture as an example of analysis of infinitesimal isometries.

Theorem 12 *There is a configuration of 12 tetrahedra with disjoint interiors such that no single tetrahedron can be moved by an infinitesimal isometry.*



Figure 5: Three of the 12 bodies

We begin with our regular tetrahedron $[\vec{a}, \vec{b}, \vec{c}, \vec{d}]$. A tetrahedral body is defined by the four vertices $72\vec{b}$, $36\vec{b}$, $72\vec{c}$, and $24(\vec{b} + \vec{d})$. Figure 5 illustrates three of the twelve bodies that are generated by the group action of A^4.

Stationary contact points on moving faces will rule out certain isometries; we determine which ones using an analysis technique that is common to closure grasps [7, 9] and assembly-sequence planning [6, 11] in robotics. Any infinitesimal rigid motion can be expressed as translational force and a rotational torque applied to the origin and can be represented as a force/torque vector with six coordinates. If we apply a force at a point p, pushing into a face with normal $\vec{n}$, then the force and torque applied to the origin is a positive multiple of the force/torque vector $\vec{f} = (-\vec{n}, -\vec{n} \times p)$, where $\times$ denotes the 3-d vector outer (or cross) product. Therefore, if we can find a set of stationary contact points whose force/torque vectors $\vec{f}_1, \ldots, \vec{f}_k$ span the six dimensional force/torque space by positive linear combinations, then those points immobilize the body—any force/torque vector will have positive projection on one of the $\vec{f}_i$.

Now, to prove that the body with vertices $72\vec{b}$, $36\vec{b}$, $72\vec{c}$, and $24(\vec{b} + \vec{d})$ cannot move rigidly, we need to find normals and contacts for its faces. Figure 6 shows the faces of this body unfolded into the plane after cutting along each of the three edges incident on vertex $36\vec{b}$. From the Cartesian coordinates of the vertices we compute four plane equations, α, β, γ and δ, with outward-pointing normals (table 2). The faces of figure 6 are labelled by the planes that contain them.

α:	$0 = ($	$1,$	$1,$	$0) \cdot q$
β:	$0 = ($	$1,$	$0,$	$1) \cdot q$
γ:	$144 = ($	$-3,$	$-3,$	$-2) \cdot q$
δ:	$-144 = ($	$2,$	$3,$	$3) \cdot q$

Table 2: Body plane eqns

As the shading in figure 6 indicates, the whole of the faces in planes α and β are in contact with faces of other bodies. In the plane γ the portion of the face bounded by $72\vec{b}$, $24(\vec{b} + \vec{d})$, and $18(\vec{b} + \vec{c} + \vec{d})$ is in contact; in the plane δ the portion bounded by $72\vec{c}$, $24(\vec{b} + \vec{c})$, and $18(\vec{b} + \vec{c} + \vec{d})$ is in contact. We choose seven contact points from the interiors of these contact regions as listed in table 3.

The interested reader can check that these vectors span the 6-d force/torque space and that $218400\vec{f}_1 + 71252\vec{f}_2 + 1548\vec{f}_3 + 43927\vec{f}_4 + 101673\vec{f}_5 + 848\vec{f}_6 +$

Plane	Contact	Force/torque vector
α :	$(35, -35, -37)$	$\vec{f}_1 = (1, 1, 0, -37, 37, -70)$
β :	$(2, -48, -2)$	$\vec{f}_2 = (1, 0, 1, 68, 132, -68)$
β :	$(66, -68, -66)$	$\vec{f}_3 = (1, 0, 1, 48, 4, -48)$
γ :	$(64, -68, -66)$	$\vec{f}_4 = (-3, -3, -2, 62, -326, 396)$
γ :	$(-16, -20, -18)$	$\vec{f}_5 = (-3, -3, -2, 14, -22, 12)$
δ :	$(-3, 1, -47)$	$\vec{f}_6 = (2, 3, 3, -144, 85, 11)$
δ :	$(-18, -16, -20)$	$\vec{f}_7 = (2, 3, 3, -12, -14, 22)$

Table 3: Contacts and force/torque vectors for the body



Figure 6: A body unfolded

$71952\vec{f}_7 = \vec{0}$. Therefore the body cannot move by any infinitesimal rigid motion.

5 A twisted triacontahedral design

In this section we form a configuration of 30 identical convex objects that cannot be taken apart with two hands using arbitrary isometries (rigid motions). The name comes from Coffin's book [1], which includes some amazing puzzles based on 30 sticks arranged around the 30 faces of a triacontahedra.

Theorem 13 *Figure 7 illustrates a configuration of 30 convex sticks with disjoint interiors that cannot be taken apart with two hands using arbitrary isometries.*

Our construction laces five copies of the twisted tetrahedron into a dodecahedron—figure 7 depicts one as solid among four wireframes on the right. These copies correspond to five subgroups of A^4 in A^5, which is the group of symmetries of a dodecahedron. By judiciously choosing separating planes, we get sticks that share two-dimensional contacts with ten neighbors.

Let $\phi = (1 + \sqrt{5})/2$ denote the golden ratio and define rotation matrices

$$F = \frac{1}{2}\begin{pmatrix} \phi & 1 & \frac{1}{\phi} \\ -1 & \frac{1}{\phi} & \phi \\ \frac{1}{\phi} & -\phi & 1 \end{pmatrix}, \; E = \frac{1}{2}\begin{pmatrix} -1 & \frac{1}{\phi} & \phi \\ \frac{1}{\phi} & -\phi & 1 \\ \phi & 1 & \frac{1}{\phi} \end{pmatrix}.$$

F and E generate A^5, the group of symmetries of a regular dodecahedron:

$$A^5 = \{F, E \mid F^5 = I, E^2 = I, (EF)^3 = I\}.$$

(Given the dodecahedron whose vertices are points in the orbit of $(1, 1, 1)$ under A^5, F rotates by $2\pi/5$ about a line through the center of a face; E rotates by π about a line through the center of an edge.) The group elements $FEFE = N$ and $EFFEF = R$ are the generators of the A^4 subgroup that we used in section 2. The group elements $\{I = F^0, F, F^2, F^3, F^4\}$ each lie in different cosets of this subgroup.

Construct sticks in a manner similar to section 2: We begin with the 60 point orbit of $p = (6, 5, -2)$ under the action of A^5. There are 30 line segments in the orbit of segment $(p; pR)$, each generated twice. We then define planes separating $(p; pR)$ from neighboring segments and intersect their half-spaces to form one stick. There are ten separating planes, which are grouped into three types in table 4. Four planes are defined to separate $(p; pR)$ from neighbors within its A^4 subgroup, four planes are defined to separate it from neighbors in an A^4 subgroup generated by R and EF, and two other bisecting planes are defined.



Figure 7: 30 objects that cannot be taken apart by two hands using arbitrary isometries

Segment	Plane eqn.
$\vec{n}_1 = (64, -79, 97)/3$	
$(pN; pRN)$	$-161/3 = \vec{n}_1 \cdot q$
$(pNR; pRNR)$	$-161/3 = \vec{n}_1 R \cdot q$
$(pNN; pRNN)$	$161/3 = \vec{n}_1 NN \cdot q$
$(pNNR; pRNNR)$	$161/3 = \vec{n}_1 NNR \cdot q$
$\vec{n}_2 = (-55 + 25\phi, 58 - 26\phi, 64 + 27\phi)/2$	
$(pEF; pREF)$	$-4 - 57\phi = \vec{n}_2 \cdot q$
$(pEFR; pREFR)$	$-4 - 57\phi = \vec{n}_2 R \cdot q$
$(pEFEF; pREFEF)$	$4 + 57\phi = \vec{n}_2 EFEF \cdot q$
$(pEFEFR; pREFEFR)$	$4 + 57\phi = \vec{n}_2 EFEFR \cdot q$
$\vec{n}_3 = (-25 - 30\phi, 30 + 36\phi, -19 + 19\phi)/2$	
$(pE; pRE)$	$0 = \vec{n}_3 \cdot q$
$(pER; pRER)$	$0 = \vec{n}_3 R \cdot q$

Table 4: Planes separating $(p; pR)$ from neighbors

These planes form non-empty, bounded, disjoint sticks. Each stick shares 2-dimensional contacts its ten neighbors, as shown in figure 8. (In this figure and in figure 8 the ends of the sticks have been clipped so that the interesting part of the construction may be seen.) We choose one point (the middle) from each contact region and form the force/torque vectors that apply to the origin when a positive force acts on each contact of each stick. We consider possible infinitesimal motions of a subset of sticks by a linear programming formulation of the analysis of the previous section: Let A be the $6 \times k$ matrix of force/torque vectors where a moving object is in contact with a stationary point. Non-degenerate basic feasible solutions of the seven linear equations $A\vec{x} = \vec{0}$ and $\sum_{1 \le i \le k} x_i = 1$ express the origin as a positive linear combination of seven force/torque vectors that span the space. If we show that this linear program is feasible for each subset of sticks, then we have shown that the sticks cannot be taken apart by any two-handed rigid motion.



Figure 8: Stick contacts

There are 2^{30} subsets to analyze; we can use a 30-bit integer to represent each subset. By manipulating these bits, we can avoid analyzing symmetric cases, complementary cases, and cases in which some moving stick has nine or ten stationary neighbors. This reduces the number of subsets to $2^{21.8}$. If we weaken the result to show that no motion to infinity is possible (rather than that no infinitesimal motion is possible) then we can reduce the number of subsets to $2^{16.8}$ by eliminating subsets with moving triangle that is linked with a stationary triangle or 4-cycle. Solving the linear programming problems for these subsets (using a modified simplex algorithm that stores only the inverse of the current

basis and not the entire tableau) takes 7 minutes on an IBM RS6000/560, including the reduction. Because these reductions also make proving program correctness more difficult, we wind up not using them on the final run of the program, which takes about 25 hours.

We should mention that, even with no pivoting, the condition numbers of all matricies that arose were bounded by 10^6 using the L_∞ norm. We ran a second check with pivoting on subsets for which the bound on the condition number became larger than 2×10^4. The actual condition numbers in these cases were less than 3.

One could implement a general polynomial-time algorithm to analyze a configuration by computing the arrangement that the hyperplanes normal to the contact force/torque vectors induce on the 5-dimensional sphere of unit force/torque vectors in 6-space. Within each cell one can define a unique *blocking graph*, with a directed edge from object A to B if contact with A prevents B from moving according to force/torque vectors in the cell. No subset can move infinitesimally iff all blocking graphs are strongly-connected. (See Wilson and Matsui [11] for more detail.) In our case, however, the 150 hyperplanes form an arrangement with 2^{34} cells, and further reductions require tricky programming.

6 Conclusion

The constructions in this paper are further evidence, if any were needed, that favorite techniques and tools of computational geometry in the plane break down in 3-space.

Many mathematical and algorithmic questions remain to be answered: Exactly how many hands are necessary to take apart any configuration of convex objects? A specific configuration? If one is told how many hands are sufficient for a specific configuration, how does one efficiently compute a motion? An entire disassembly sequence?

Acknowledgements

We thank Simon Kahan and Dale Rolfsen for helping to debug proofs that one of five objects can translate and thank Randy Wilson for his paper on the polynomial-time disassembly algorithm and other references to the literature on mechanical assembly planning.

References

[1] S. T. Coffin. *The Puzzling World of Polyhedral Dissections*. Oxford University Press, 1991.

[2] R. Dawson. On removing a ball without disturbing the others. *Math. Mag.*, 57(1):27–30, Jan. 1984.

[3] N. G. de Bruijn. *Nieuw Archief voor Wiskunde*, 2:67, 1954. Problems 17 and 18. Answers in *Wiskundige Opgaven met de oplossingen*, 20:19–20, 1955.

[4] L. Fejes Toth and A. Heppes. Uber stabile Körpersysteme. *Compositio Mathematica*, 15(2):119–126, 1963.

[5] L. J. Guibas and F. F. Yao. On translating a set of rectangles. In *Proc. 12th Ann. ACM STOC*, pages 154–160, 1980.

[6] L. S. H. d. Mello. *Computer-Aided Mechanical Assembly Planning*. Kluwer Academic Publishers, Boston, 1991.

[7] B. Mishra, J. T. Schwartz, and M. Sharir. On the existence and synthesis of multifinger positive grips. *Algorithmica*, 2(541–558), 1987.

[8] B. K. Natarajan. On planning assemblies. In *Proc. 4th Ann. ACM Symp. Comp. Geom.*, pages 299–308, 1988.

[9] J. Pertin-Troccaz. Grasping: A state of the art. In O. Khatib, J. J. Craig, and T. Lozano-Perez, editors, *The Robotics Review 1*, pages 71–98. MIT Press, 1989.

[10] F. P. Preparata. Planar point location revisited. *Int. J. Found. Comp. Sci.*, 1(1):71–86, 1990.

[11] R. H. Wilson and T. Matsui. Partitioning an assembly for infinitesimal motions in translation and rotation. In *IEEE International Conference on Intellegent Robots and Systems*, pages 1311–1318, 1992.